

A new iterative algorithm on the time-fractional Fisher equation: Residual power series method

Maysaa' Mohamed Al Qurashi¹, Zeliha Korpınar², Dumitru Baleanu^{3,4} and Mustafa Inc⁵

Abstract

In this article, the residual power series method is used to solve time-fractional Fisher equation. The residual power series method gets Maclaurin expansion of the solution. The solutions of present equation are computed in the shape of quickly convergent series with quickly calculable fundamentals using mathematica software package. Explanation of the method is given by graphical consequences and series solutions are made use of to represent our solution. The found consequences show that technique is a power and efficient method in conviction of solution time-fractional Fisher equation.

Keywords

Residual power series method, time-fractional Fisher equation, series solution

Date received: 4 January 2017; accepted: 14 May 2017

Academic Editor: Xiao-Jun Yang

Introduction

Important care has been assigned to the work of the fractional calculus during the last few decades and its numerous utilizations in the physics, regular variation in biophysics, thermodynamics, blood flow phenomena, viscoelasticity, electrical circuits, aerodynamics, astrophysics, biology, control theory, and so on.^{1–4} However, fractional derivatives supply an important implement for the definition of recollection and hereditary characteristics of different necessities and treatment. This is the fundamental advantage of fractional differential equations in return usual integer order problems.

Recently, there has been a significant analytical improvement in fractional differential equations and its applications. In the work by Yang et al.,⁵ researchers applied the local fractional derivative operator for obtaining the non-differential solution for diffusion equation in fractal heat transfer; in the work by Gao et al.,⁶ the exact solution for the local fractional diffusion equation in fractal one-dimensional space is

obtained; in the work by Yang et al.,⁷ the analytical solutions of the sub-diffusion and wave equations are obtained by utilizing the local fractional variational iteration method; in the work by Gao and Yang,⁸ the local fractional Euler's method is applied to obtain numerical solution for the local fractional heat-relaxation equation; in the work by Yang et al.,⁹ authors analyzed the exact traveling wave solutions for local fractional Korteweg–de Vries (KdV) equation; in

¹Department of Mathematics, King Saud University, Riyadh, Saudi Arabia

²Department of Administration, Faculty of Economic and Administrative Sciences, Muş Alparslan University, Muş, Turkey

³Department of Mathematics and Computer Sciences, Cankaya University, Ankara, Turkey

⁴Institute of Space Sciences, Măgurele, Romania

⁵Department of Mathematics, Faculty of Science, Firat University, Elazığ, Turkey

Corresponding author:

Dumitru Baleanu, Department of Mathematics and Computer Sciences, Cankaya University, Ankara, 06790, Turkey.
Email: dumitru@cankaya.edu.tr



the work by Yang et al.,¹⁰ the exact traveling wave solutions for the local fractional two-dimensional Burgers-type equations are investigated, and in the work by Zhang et al.,¹¹ researchers utilized the series expansion method within local fractional derivative to obtain the solutions of homogeneous and non-homogeneous transport equations, respectively. For some articles and books on fractional differential and fractional calculus equations, see Baleanu and colleagues^{12–14} and the references therein.

In this study, we apply residual power series method (RPSM) to find powerful series solution for a nonlinear problem. The new method supplies the solution in the shape of a convergence series. A repeated algorithm is constituted for the designation of the infinite series solution. The RPSM was developed as an efficient method for fuzzy differential equations.¹⁵ It has been successfully put into practice to handle the approximate solution for generalized Lane–Emden equation,¹⁶ the solution of compound and non-compound fractional differential equations,¹⁷ predicting and representing the profusion of results to boundary value issues of fractional order,¹⁸ constructing and forecasting the solitary arrangement solutions for nonlinear time-fractional advance partial differential equations,¹⁹ the comparative solution of the nonlinear fractional KdV–Burgers equation,²⁰ the comparative solutions of fractional population diffusion model,²¹ and the numerical solutions of linear non-homogeneous partial differential equations of fractional order.²² The introduced method constitutes a comparative solution in the shape of a polynomial. Contrary to Taylor series method, RPSM needs small calculational condition with high certainty and less time at older orders. RPSM is a preference process for getting consistent Maclaurin series solution of problems.

The base purpose of our work is to present practice of RPSM in the touch of the Caputo fractional differential to examine and establish an approximate solution of the space-time-fractional order Fisher equation²³

$$D_t^\alpha u(x, t) = D_{xx}u(x, t) + 6u(x, t)(1 - u(x, t)), \quad (1)$$

$$x \in R, t > 0, 0 < \alpha \leq 1$$

In above equation, $u(x, t)$ is estimated to be a function of space and time, which equipment that $u(x, t)$ is disappearing for $t < 0$ and $x < 0$ and this function is considered to be analytic on $t > 0$. Moreover, the function $f(x)$ is considered to be analytic on $x > 0$.

Time-fractional Fisher equation can be readily solved by many methods. In the work by Rida et al.,²⁴ an accurate algorithm for the solution of special conditions is developed and differential transform method is used. Khan et al.²⁵ used homotopy perturbation approach in order to find the comparative solutions of time-fractional Fisher equation. Sungu and Demir²³

applied generalized differential transform and finite difference method for numerical solutions of the equation. Baranwal et al.²⁶ and Merdan²⁷ used fractional variational iteration method for series solutions of proposed equation.

In section “Some necessary definitions and results from fractional calculus” of this work, some preliminary results related to the Caputo derivative and the fractional power series (PS) are described. In section “Solution of the time-fractional Fisher equation with RPSM algorithm,” base opinion of the RPSM is constituted to construct the solution of the time-fractional Fisher equation (1). In section “Graphical results of the RPSM,” some graphical consequences are included to demonstrate the reliability and efficiency of the method. Moreover, consequences are introduced in section “Final remarks.”

Some necessary definitions and results from fractional calculus

Below, we first illustrate the essential descriptions and various features for the fractional algebra theory.²

Definition 1. From El-Ajou et al.,^{17,20} the fractional integral operator Riemann–Liouville with order α ($\alpha \geq 0$) is given by

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad \alpha > 0, x > 0 \quad (2)$$

$$J^0 f(x) = f(x)$$

Definition 2. The Caputo fractional derivative with order α is given by

$$D^\alpha f(x) = J^{m-\alpha} D^m f(x) = \frac{1}{\Gamma(m-\alpha)} \int_0^x (x-t)^{m-\alpha-1} \frac{d^m}{dt^m} f(t) dt, \quad m-1 < \alpha \leq m, x > 0 \quad (3)$$

where D^m is the classic differential operator with order m .^{17,20}

By the Caputo derivative, we get

$$D^\alpha x^\beta = 0, \quad \beta < \alpha$$

$$D^\alpha x^\beta = \frac{\Gamma(\beta+1)}{\Gamma(\beta+1-\alpha)} x^{\beta-\alpha}, \quad \beta \geq \alpha$$

Definition 3. For n to be the smallest integer that exceeds α , the Caputo time-fractional differential operator of order α of $u(x, t)$ is defined as^{17,20}

$$D_t^\alpha u(x, t) = \frac{\partial^\alpha u(x, t)}{\partial t^\alpha} = \frac{1}{\Gamma(n - \alpha)}$$

$$\int_0^t (t - \tau)^{n-\alpha-1} \frac{\partial^n u(x, \tau)}{\partial t^n} d\tau, \quad n - 1 < \alpha < n \quad (4)$$

$$D_t^n u(x, t) = \frac{\partial^n u(x, t)}{\partial t^n}, \quad n \in \mathbb{N}$$

and the space-time-fractional differential with order β of $u(x, t)$ is defined by

$$D_x^\beta u(x, t) = \frac{\partial^\beta u(x, t)}{\partial x^\beta} = \frac{1}{\Gamma(n - \beta)}$$

$$\int_0^x (x - \tau)^{n-\beta-1} \frac{\partial^n u(\tau, t)}{\partial t^n} d\tau, \quad n - 1 < \beta < n \quad (5)$$

$$D_x^n u(x, t) = \frac{\partial^n u(x, t)}{\partial x^n}, \quad n \in \mathbb{N}$$

Definition 4. A PS expanding of the manner

$$\sum_{m=0}^{\infty} c_m (t - t_0)^{m\alpha} = c_0 + c_1 (t - t_0)^\alpha$$

$$+ c_2 (t - t_0)^{2\alpha} + \dots, \quad 0 \leq m - 1 < \alpha \leq m, \quad t \geq t_0$$

is named fractional PS at $t = t_0$.¹³

Definition 5. A PS of the form

$$\sum_{m=0}^{\infty} f_m(x) (t - t_0)^{m\alpha} = f_0(x) + f_1(x) (t - t_0)^\alpha$$

$$+ f_2(x) (t - t_0)^{2\alpha} + \dots, \quad 0 \leq m - 1 < \alpha \leq m, \quad t \geq t_0 \quad (6)$$

is named fractional PS at $t = t_0$.¹³

Theorem 1. Only $u(x, t)$ gives a polynomial fractional PS representing at point $t = t_0$ of the shape (see El-Ajou et al.²⁰ for proof)

$$u(x, t) = \sum_{m=0}^{\infty} f_m(x) (t - t_0)^{m\alpha}, \quad (7)$$

$$0 \leq m - 1 < \alpha \leq m, \quad x \in I, \quad t_0 \leq t < t_0 + R$$

If $D_t^{m\alpha} u(x, t)$ are continuous on $I \times (t_0, t_0 + R)$, in this case, coefficients $f_m(x)$ are given as

$$f_m(x) = \frac{D_t^{m\alpha} u(x, t_0)}{\Gamma(m\alpha + 1)}, \quad m = \overline{0, \infty}$$

where $D_t^{m\alpha} = \partial^{m\alpha} / \partial t^{m\alpha} = (\partial^\alpha / \partial t^\alpha) \cdot (\partial^\alpha / \partial t^\alpha) \dots (\partial^\alpha / \partial t^\alpha)$ (m times) and $R = \min_{c \in I} R_c$, in which R_c is the effect domain of convergency of the fractional PS $\sum_{m=0}^{\infty} f_m(c) (t - t_0)^{m\alpha}$.

Result 1. The fractional PS expanded of $u(x, t)$ at point t_0 should be of the shape

$$u(x, t) = \sum_{m=0}^{\infty} \frac{D_t^{m\alpha} u(x, t_0)}{\Gamma(m\alpha + 1)} (t - t_0)^{m\alpha}, \quad (8)$$

$$0 \leq m - 1 < \alpha \leq m, \quad x \in I, \quad t_0 \leq t < t_0 + R$$

which denotes a generalized Taylor's series expression. To particularize, if one sets $\alpha = 1$ in equation (8), in this case, the classical Taylor's series expression

$$u(x, t) = \sum_{m=0}^{\infty} \frac{\partial^m u(x, t_0)}{\partial t^m} \frac{(t - t_0)^m}{m!}, \quad x \in I, \quad t_0 \leq t < t_0 + R$$

is obtained.²⁰

Solution of the time-fractional Fisher equation with RPSM algorithm

First, we study the time-fractional Fisher equation

$$D_t^\alpha u(x, t) = D_{xx} u(x, t) + 6u(x, t)(1 - u(x, t)), \quad (9)$$

$$x \in R, \quad t > 0, \quad 0 < \alpha \leq 1$$

by the initial condition

$$u(x, 0) = \frac{1}{(1 + e^x)^2} \quad (10)$$

Therefore, we study procedure of the RPSM to ascertain series solution for our time-fractional Fisher equation dependent on given initial conditions by replacing its fractional PS expanded with its truncated residual function.

The RPSM proposes the solution for equations (9) and (10) with a fractional PS at point $t = 0$.¹⁵ Theorize that the solution selects the expansion shape

$$u(x, t) = \sum_{n=0}^{\infty} f_n(x) \frac{t^{n\alpha}}{\Gamma(1 + n\alpha)} \quad 0 < \alpha \leq 1, \quad x \in I, \quad 0 \leq t < R \quad (11)$$

Next, we let $u_k(x, t)$ to refer k truncated series of $u(x, t)$

$$u_k(x, t) = \sum_{n=0}^k f_n(x) \frac{t^{n\alpha}}{\Gamma(1 + n\alpha)}, \quad 0 < \alpha \leq 1, \quad x \in I, \quad 0 \leq t < R \quad (12)$$

where $u_0(x, t) = f_0(x) = u(x, 0) = f(x)$.

Besides, equation (12) may be reported by

$$u_k(x, t) = f(x) + \sum_{n=1}^k f_n(x) \frac{t^{n\alpha}}{\Gamma(1 + n\alpha)}, \quad (13)$$

$$0 < \alpha \leq 1, \quad 0 \leq t < R, \quad x \in I, \quad k = \overline{1, \infty}$$

At first, to find the value of coefficients $f_n(x)$, $n = 1, 2, 3, \dots, k$ in series expanded of equation (13), we define residual function Res for equation (1) by

$$Res(x, t) = \frac{\partial^\alpha u(x, t)}{\partial t^\alpha} - \frac{\partial^2 u(x, t)}{\partial x^2} - 6u(x, t)(1 - u(x, t))$$

and the k th residual function Res_k as follows

$$Res_k(x, t) = \frac{\partial^\alpha u_k(x, t)}{\partial t^\alpha} - \frac{\partial^2 u_k(x, t)}{\partial x^2} - 6u_k(x, t)(1 - u_k(x, t)), \quad k = 1, 2, 3, \dots \quad (14)$$

As in Abu Arqub and colleagues,¹⁵⁻¹⁸ it is clear that $Res(x, t) = 0$ and $\lim_{k \rightarrow \infty} Res_k(x, t) = Res(x, t)$ for each $x \in I$ and $t \geq 0$.

Then, $D_t^\alpha Res(x, t) = 0$, fractional derivative of a stationary in the Caputo's idea is zero, and the fractional derivative D_t^α of $Res(x, t)$ and $Res_k(x, t)$ is pairing at $t = 0$ with each $r = \overline{0, k}$. To give residual PS algorithm, first, we replace the k th truncated series of $u(x, t)$ with equation (9). Second, we find the fractional derivative expression $D_t^{(k-1)\alpha}$ of both $Res_{u, k}(x, t)$ and $k = \overline{1, \infty}$, and, finally, we can solve found system

$$D_t^{(k-1)\alpha} Res_{u, k}(x, 0) = 0, \quad 0 < \alpha \leq 1, x \in I, k = \overline{1, \infty} \quad (15)$$

to obtain the needed coefficients $f_n(x)$ for $n = \overline{1, k}$ in equation (13).

Hence, to determine $f_1(x)$, we write $k = 1$ in equation (14)

$$Res_1(x, t) = \frac{\partial^\alpha u_1(x, t)}{\partial t^\alpha} - \frac{\partial^2 u_1(x, t)}{\partial x^2} - 6u_1(x, t)(1 - u_1(x, t)) \quad (16)$$

where

$$u_1(x, t) = \frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + f(x)$$

for

$$u(x, 0) = f_0(x) = f(x) = u(x, 0) = \frac{1}{(1 + e^x)^2}$$

Therefore

$$Res_1(x, t) = f_1(x) - f''(x) - \frac{t^\alpha}{\Gamma(1 + \alpha)} f''_1(x) - 6 \left(\frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + f(x) \right) \left(1 - \left(\frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + f(x) \right) \right)$$

From equation (15), we deduce that $Res_1(x, 0) = 0$ and thus

$$f_1(x) = \frac{10e^x}{(1 + e^x)^3} \quad (17)$$

Therefore

$$u_1(x, t) = \frac{10e^x}{(1 + e^x)^3} \frac{t^\alpha}{\Gamma(1 + \alpha)} + \frac{1}{(1 + e^x)^2} \quad (18)$$

Likewise, to determine the shape of the second unnamed coefficient $f_2(x)$, we write $k = 2$ in equation (14)

$$Res_2(x, t) = \frac{\partial^\alpha u_2(x, t)}{\partial t^\alpha} - \frac{\partial^2 u_2(x, t)}{\partial x^2} - 6u_2(x, t)(1 - u_2(x, t))$$

where

$$u_2(x, t) = f(x) + \frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} f_2(x)$$

Therefore

$$Res_2(x, t) = f_1(x) + \frac{t^\alpha}{\Gamma(1 + \alpha)} f_2(x) - f''(x) - \frac{t^\alpha}{\Gamma(1 + \alpha)} f''_1(x) - \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} f''_2(x) - 6 \left(f(x) + \frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} f_2(x) \right) \left(1 - f(x) - \frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) - \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} f_2(x) \right)$$

From equation (15), we deduce that $D_t^\alpha Res_2(x, 0) = 0$ and thus

$$f_2(x) = \frac{50e^x(-1 + 2e^x)}{(1 + e^x)^4} \quad (19)$$

Therefore

$$u_2(x, t) = \frac{1}{(1 + e^x)^2} + \frac{10e^x}{(1 + e^x)^3} \frac{t^\alpha}{\Gamma(1 + \alpha)} + \frac{50e^x(-1 + 2e^x)}{(1 + e^x)^4} \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} \quad (20)$$

Similarly, to determine $f_3(x)$, we write $k = 3$ in equation (14)

$$Res_3(x, t) = \frac{\partial^\alpha u_3(x, t)}{\partial t^\alpha} - \frac{\partial^2 u_3(x, t)}{\partial x^2} - 6u_3(x, t)(1 - u_3(x, t))$$

where

$$u_3(x, t) = f(x) + \frac{t^\alpha}{\Gamma(1 + \alpha)} f_1(x) + \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} f_2(x) + \frac{t^{3\alpha}}{\Gamma(1 + 3\alpha)} f_3(x)$$

Therefore

$$\begin{aligned}
Res_3(x, t) = & f_1(x) + \frac{t^\alpha}{\Gamma(1+\alpha)}f_2(x) + \frac{t^{2\alpha}}{\Gamma(1+2\alpha)}f_3(x) \\
& - \left(f''(x) + \frac{t^\alpha}{\Gamma(1+\alpha)}f''_1(x) + \frac{t^{2\alpha}}{\Gamma(1+2\alpha)}f''_2(x) + \frac{t^{3\alpha}}{\Gamma(1+3\alpha)}f''_3(x) \right) \\
& - 6 \left(f(x) + \frac{t^\alpha}{\Gamma(1+\alpha)}f_1(x) + \frac{t^{2\alpha}}{\Gamma(1+2\alpha)}f_2(x) + \frac{t^{3\alpha}}{\Gamma(1+3\alpha)}f_3(x) \right) \\
& \left(1 - \left(f(x) + \frac{t^\alpha}{\Gamma(1+\alpha)}f_1(x) + \frac{t^{2\alpha}}{\Gamma(1+2\alpha)}f_2(x) + \frac{t^{3\alpha}}{\Gamma(1+3\alpha)}f_3(x) \right) \right)
\end{aligned}$$

From equation (15), we deduce that $D_t^{2\alpha}Res_3(x, 0) = 0$ and thus

$$f_3(x) = \frac{125e^x(1 - 7e^x + 4e^{2x})}{(1 + e^x)^5} \quad (21)$$

Therefore

$$\begin{aligned}
u_3(x, t) = & \frac{10e^x}{(1 + e^x)^3} \frac{t^\alpha}{\Gamma(1+\alpha)} + \frac{1}{(1 + e^x)^2} \\
& + \frac{50e^x(-1 + 2e^x)}{(1 + e^x)^4} \frac{t^{2\alpha}}{\Gamma(1+2\alpha)} \\
& + \frac{125e^x(1 - 7e^x + 4e^{2x})}{(1 + e^x)^5} \frac{t^{3\alpha}}{\Gamma(1+3\alpha)}
\end{aligned} \quad (22)$$

Likewise, employing the similar operation for $k = 4$ taking describe the shape of $f_0(x), f_1(x), f_2(x), f_3(x)$, particularly, will moderate after easily computations to the next shape of $f_4(x)$

$$f_4(x) = \frac{625e^x(-1 + 18e^x - 33e^{2x} + 8e^{3x})}{3(1 + e^x)^6} \quad (23)$$

Therefore

$$\begin{aligned}
u_4(x, t) = & \frac{1}{(1 + e^x)^2} + \frac{10e^x}{(1 + e^x)^3} \frac{t^\alpha}{\Gamma(1+\alpha)} \\
& + \frac{50e^x(-1 + 2e^x)}{(1 + e^x)^4} \frac{t^{2\alpha}}{\Gamma(1+2\alpha)} + \frac{125e^x(1 - 7e^x + 4e^{2x})}{(1 + e^x)^5} \\
& \frac{t^{3\alpha}}{\Gamma(1+3\alpha)} + \frac{625e^x(-1 + 18e^x - 33e^{2x} + 8e^{3x})}{3(1 + e^x)^6} \frac{t^{4\alpha}}{\Gamma(1+4\alpha)}
\end{aligned} \quad (24)$$

To give a deficit overview of the content of our work, in the next section, by the above recurrent connections, we can demonstrate some graphical consequences of equations (1) and (2) in Figures 1–3.




Figure 1. The 3D graphic for the exact solution $u(x, t)$ and the $u_4(x, t)$ approximate solution of the time-fractional Fisher equation: (a) $u_4(x, t)$ when $\alpha = 0.5$, (b) $u_4(x, t)$ when $\alpha = 0.75$, (c) $u_4(x, t)$ when $\alpha = 0.9$, and (d) $u(x, t)$ when $\alpha = 1$.




Figure 2. $u_k(x, t)$ solution of the time-fractional Fisher equation when $k = 1, 2, 3, 4$ versus its exact solution: (a) $\alpha = 0.9$, $t = 0.4$ and (b) $\alpha = 0.5$, $t = 0.1$.




Figure 3. $u_4(x, t)$ solution of the time-fractional Fisher equation when $\alpha = 0.5, 0.6, 0.7, 0.8, 0.9, 1$ ($t = 0.1$ and $t = 0.3$).

Graphical results of the RPSM

Following our considerations in the previous section, we can form graphical consequences of equation

$$D_t^\alpha u(x, t) = D_{xx}u(x, t) + 6u(x, t)(1 - u(x, t)),$$

$$t > 0, x \in \mathbb{R}, 0 < \alpha \leq 1$$

with the initial condition

$$u(x, 0) = \frac{1}{(1 + e^x)^2}$$

$u(x, t) = 1/(1 + e^{x-5t})^2$ is the exact solution of this equation.²⁴

In Figure 2, we plot the RPS (residual power series) approximate solution $u_k(x, t)$ for $k = 1, 2, 3, 4$ which are closing the line $y = 0$ as the amount of iterations enlarge. This figure clearly shows that the absolute error is smaller as the amount of k enlarges. It is clear that the value of k th truncated series $u_k(x, t)$ affects the RPS approximate solutions.

In Figure 3, we plot the RPS approximate solution $u_4(x, t)$ for $\alpha = 0.5 - 1$ which is closing the exact

Table 1. Comparison among approximate solutions u_{RPSM} , u_{FVM} , and u_{HPM} ($x = 0.5$).

$\alpha = 0.75$			
t	u_{RPSM}	u_{FVM}	u_{HPM}
0.01	0.175966	0.0242651	0.146808
0.05	0.277218	0.0818379	0.157343
0.1	0.406898	0.139343	0.168312
0.15	0.54626	0.186505	0.178529
0.2	0.695654	0.224693	0.188419

solution as the number of α increases. This figure clear shows that the convergency of the approximate solutions to the exact solution related to the order of the solution and the exact error is smaller as the order of the solution increases.

In Tables 1–3, we constitute table of numerical solutions $u_k(x, t)$ for $k = 3$. Comparison among approximate solutions with known results is made. These results are obtained using RPSM, homotopy perturbation method,²⁵ and fractional variational iteration method.²⁷

Table 2. Comparison among approximate solutions u_{RPSM} , u_{FVM} , and u_{HPM} ($x = 0.5$).

$\alpha = 0.5$			
t	u_{RPSM}	u_{FVM}	u_{HPM}
0.01	0.281524	0.125003	0.157224
0.05	0.563205	0.316523	0.177994
0.1	0.856396	0.450853	0.195594
0.15	1.13039	0.528601	0.210357
0.2	1.39515	0.567751	0.22369

Table 3. Comparison between approximate solution u_{RPSM} and exact solution ($x = 0.5$).

$\alpha = 1$			
t	u_{RPSM}	u_{Exact}	$ u_{Exact} - u_{RPSM} $
0.01	0.151602	0.151602	3.46783×10^{-8}
0.05	0.191713	0.191689	2.34838×10^{-5}
0.1	0.250483	0.25	4.82771×10^{-4}
0.15	0.318677	0.316042	2.63503×10^{-3}
0.2	0.395971	0.387456	8.51568×10^{-3}

Tables 1–3 clarify the convergency of the approximate solutions to the exact solution and exact error is smaller as the value of the t decreases.

Final remarks

The fundamental objective of this article is to introduce an algorithmic form and implement a new analytical repeated algorithm derived from the RPS to find numerical solutions for nonlinear time-fractional Fisher equation. The approximate solution is given in the shape of PS. Graphical and numerical consequences are introduced to illustrate the solutions. The consequences found using the RPSM are very active and appropriate in general conditions with less calculational work and time. It has been established that the structure of this RPSM possesses a very fast convergent series with easily calculable components using symbolic calculation software. The article stressed our notion that the introduced process can be applied as an instead to get analytic solutions of different kinds of fractional linear and nonlinear partial differential equations practiced in mathematics, physics, and engineering.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was funded by the International Scientific Partnership Program ISPP at King Saud University (ISPP# 63).

References

- Kilbas AA, Srivastava HM and Trujillo JJ. *Theory and applications of fractional differential equations*. Amsterdam: Elsevier, 2006.
- Podlubny I. *Fractional differential equation*. San Diego, CA: Academic Press, 1999.
- Sabatier J, Agrawal OP and Machado JAT (eds). *Advances in fractional calculus: theoretical developments and applications in physics and engineering*. Dordrecht: Springer, 2007.
- Samko SG, Kilbas AA and Marichev OI. *Fractional integrals and derivatives: theory and applications*. Lausanne: Gordon & Breach, 1993.
- Yang X-J, Tenreiro Machado JA, Baleanu D, et al. A new numerical technique for local fractional diffusion equation in fractal heat transfer. *J Nonlinear Sci Appl* 2016; 9: 5621–5628.
- Gao F, Srivastava HM, Gao Y-N, et al. A coupling method involving the Sumudu transform and the variational iteration method for a class of local fractional diffusion equations. *J Nonlinear Sci Appl* 2016; 9: 5830–5835.
- Yang X-J, Baleanu D, Khan Y, et al. Local fractional variational iteration method for diffusion and wave equations on cantor sets. *Roman J Phys* 2014; 59: 36–48.
- Gao F and Yang X-J. Local fractional Euler's method for the steady heat-conduction problem. *Thermal Sci* 2016; 20: 735–738.
- Yang X-J, Tenreiro Machado JA, Baleanu D, et al. On exact traveling-wave solutions for local fractional Korteweg-de Vries equation. *Chaos* 2016; 26: 084312.
- Yang X-J, Gao F and Srivastava HM. Exact travelling wave solutions for the local fractional two-dimensional Burgers-type equations. *Comput Math Appl* 2017; 73: 203–210.
- Zhang Y, Baleanu D and Yang X-J. New solutions of the transport equations in porous media within local fractional derivative. *Proc Rom Acad* 2016; 17: 230–236.
- Yang X-J, Baleanu D and Zhong W-P. Approximate solutions for diffusion equations on cantor space-time. *Proc Rom Acad* 2013; 14: 127–133.
- Magin R, Feng X and Baleanu D. Solving the fractional order Bloch equation. *Concept Magnetic Res* 2009; 34A: 16–23.
- Baleanu D. New applications of fractional variational principles. *Rep Math Phys* 2008; 61: 199–206.
- Abu Arqub O. Series solution of fuzzy differential equations under strongly generalized differentiability. *J Adv Res Appl Math* 2013; 5: 31–52.
- Abu Arqub O, El-Ajou A, Bataineh A, et al. A representation of the exact solution of generalized Lane Emden

- equations using a new analytical method. *Abstr Appl Anal* 2013; 2013: 378593.
17. El-Ajou A, Abu Arqub O, Al Zhour Z, et al. New results on fractional power series: theories and applications. *Entropy* 2013; 15: 5305–5323.
 18. Abu Arqub O, El-Ajou A, Al Zhour Z, et al. Multiple solutions of nonlinear boundary value problems of fractional order: a new analytic iterative technique. *Entropy* 2014; 16: 471–493.
 19. Abu Arqub O, El-Ajou A and Momani S. Constructing and predicting solitary pattern solutions for nonlinear time-fractional dispersive partial differential equations. *J Comput Phys* 2015; 293: 385–399.
 20. El-Ajou A, Abu Arqub O and Momani S. Approximate analytical solution of the nonlinear fractional KdV–Burgers equation: a new iterative algorithm. *J Comput Phys* 2015; 293: 81–95.
 21. Alquran M, Al-Khaled K and Chattopadhyay J. Analytical solutions of fractional population diffusion model: residual power series. *Nonlin Stud* 2015; 22: 31–39.
 22. El-Ajou A, Abu Arqub O, Momani S, et al. A novel expansion iterative method for solving linear partial differential equations of fractional order. *Appl Math Comput* 2015; 257: 119–133.
 23. Sungu IC and Demir H. New approach and solution technique to solve time fractional nonlinear reaction-diffusion equations. *Math Probl Eng* 2015; 2015: 457013.
 24. Rida SZ, El-Sayed AMA and Arafa AAM. On the solutions of time-fractional reaction–diffusion equations. *Commun Nonlinear Sci* 2010; 15: 3847–3854.
 25. Khan NA, Ayaz F, Jin L, et al. On approximate solutions for the time-fractional reaction-diffusion equation of Fisher type. *Int J Phys Sci* 2011; 6: 2483–2496.
 26. Baranwal VK, Pandey RK, Tripathi MP, et al. An analytic algorithm for time fractional nonlinear reaction–diffusion equation based on a new iterative method. *Commun Nonlinear Sci* 2012; 17: 3906–3921.
 27. Merdan M. Solutions of time-fractional reaction–diffusion equation with modified Riemann–Liouville derivative. *Int J Phys Sci* 2012; 7: 2317–2326.