

## An Investigation of Anticholinesterase and Anticancer Effects of *Verbascum insulare* Boiss. Et Heldr. Extracts Growing in Muş Region

Yusuf ALAN<sup>1</sup>, Züleyha ALMAZ<sup>1\*</sup>



<sup>1</sup>Department of Molecular Biology and Genetics, Faculty of Arts and Sciences, Mus Alparslan University, Mus, 49250, Turkey  
(ORCID: [0000-0003-0007-0212](https://orcid.org/0000-0003-0007-0212)) (ORCID: [0000-0002-4532-4311](https://orcid.org/0000-0002-4532-4311))

**Keywords:** *Verbascum insulare* Boiss. Et Heldr., Anticancer, Anticholinesterase.

### Abstract

Since the existence of humanity, plants have been used in many treatment methods. Members of the genus *Verbascum*, a member of the Scrophulariaceae family, also known as mullein, have recently received great attention due to their remarkable biological activities and have been evaluated in traditional therapeutic uses against various ailments. In this study, the anticholinesterase and anticancer activities of *Verbascum* member *Verbascum insulare* Boiss. Et Heldr. collected from Muş region were investigated. *V.insulare* leaf ethanol (L-EtOH), leaf purified water (L-PW), root ethanol (R-EtOH), and root purified water (R-PW) extracts were obtained. The anticancer activity of the extracts against HT-29, MCF-7, and L-929 cell lines was examined using the 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) test, and the anticholinesterase activity of the extracts was determined. It was found that the ethanol extract showed higher anticancer activity against cancer cell lines than the pure water extract and had more effective anticholinergic influences. It is aimed to shed light on the biological effects of *Verbascum insulare* Boiss. Et Heldr, which is thought to be a natural source of bioactive compounds for the development of phytopharmaceuticals targeting oxidative stress-related diseases such as cancer and neurological diseases. No study has been found in the literature on the anticancer and anticholinesterase activity of this plant yet. Therefore, the results obtained from this study will contribute significantly to the development of herbal medicine in the future.

### 1. Introduction

Nowadays, plants are of great interest all over the world as an effective therapeutic agent for the treatment and prevention of diseases. In addition, plants, which form an important component of foodstuffs, have formed the basis of traditional medicine throughout history. It has been determined that there are approximately 420,000 plant species on earth, and more than 35,000 of these plants are used for medicinal purposes [1]. *Verbascum* genus, a member of the Scrophulariaceae family, also known as “king’s candle or mullein” in the flora of Turkey, has 233 species, 185 of which are endemic,

and more than 100 hybrids [2]. In traditional Turkish medicine, all aerial parts of *Verbascum* L. (Scrophulariaceae) species are used as a drying agent for wounds such as inflammatory skin disorders. In addition, some species of the plant are reported to have expectorant, mucolytic and sedative properties used in the treatment of respiratory ailments such as bronchitis, tuberculosis, and asthma [3]. *Verbascum nubicum* Murb species showed significant hepatoprotective, anti-inflammatory, and antioxidant effects[4]. *Verbascum bombyciferum* Boiss inhibited several enzymes, including acetylcholinesterase and butyrylcholinesterase[5]. In a study investigating the

\*Corresponding author: [z.turkoglu@alparslan.edu.tr](mailto:z.turkoglu@alparslan.edu.tr)

Received: 27.12.2021, Accepted: 29.04.2022

antiproliferative effect of different extracts of *Verbascum sinaiticum* against Hep-2, MCF-7, and Vero cell lines, it was determined that the ethanol extract of the plant exhibited high antiproliferative potential against the tested cell lines [6]. In a study of the phytochemical examination of *Verbascum thapsus*, it was determined that luteolin and 3-O-fucopyranosylsaikogenin F compounds induced apoptosis of A549 lung cancer cells and showed promising antiproliferative activities as a result [7]. Phenolic content of *Verbascum insulare* Boiss. Et Heldr., antimicrobial, antioxidant, and DNA protective activities of different extracts were determined. It was found that ethanolic extract indicated higher antioxidant and antifungal activity and increased DNA stability than pure aqueous extract [2].

Due to the increase in the number of older adults, the prevalence of diseases such as cancer and neurodegenerative disorders has increased greatly in the world in recent years. Alzheimer's disease (AD) is the most common neurodegenerative disease that progresses over time and is very difficult to diagnose, seen in middle-aged and elderly individuals. This disease is characterized by early progressive loss of neuronal cells [8]. Cancer is characterized by inappropriate cell proliferation and resistance to cell death. The common causative factor of AD and cancer is thought to be aging. It is possible to associate cancer and Alzheimer's disease with various mechanisms, and one of these mechanisms is that both diseases increase oxidative stress and thus tumor or neuronal cells become vulnerable [9]. It is known that oxidative stress increases colon and breast cancer types [10]. Consequently, new, safe and effective therapeutic approaches are needed to control these diseases. Considering the above, many *Verbascum* species can be considered promising raw materials for the design of new pharmaceutical products. While many *Verbascum* species have been studied, more research is needed to provide scientifically validated information, particularly on poorly studied species. Our aim with this research article is to investigate the anticholinesterase and anticancer activities of the medicinally important plant *V.insulare* and determine whether it can be used in the pharmaceutical industry.

## 2. Material and Method

### 2.1. The Extraction and Collection of Plant Samples

*V.insulare* plant was collected during the vegetation period in 2018 from Guzeltepe village of Mus province. The identification of the collected plant samples according to the Flora of Turkey was done by Murat Kurşat (Bitlis, Turkey). The plant samples were turned into herbarium material (Y. Alan: 4913) and stored in Muş Alparslan University, Central Research Laboratories Application and Research Center. The leaves and roots of the plant were removed and left to dry in the shade. *V.insulare* leaf ethanol (L-EtOH), leaf purified water (L-PW), root ethanol (R-EtOH), and root purified water (R-PW) extracts were prepared according to our previous study [2].

### 2.2. Cholinesterase Inhibition Assay

Acetylcholinesterase (AChE) from Electric eel (*Electrophorus electricus*) and butyrylcholinesterase (BChE) from equine serum were obtained from Sigma- Aldrich (St. Louise, MO). The inhibitory effects of *V. insulare* extracts against AChE and BChE were measured using a slight modification of Ellman's spectrophotometric method [11] as described previously [12] and using commercially available galantamine as the reference compound. Acetylthiocholine iodide (AChI), butyrylthiocholine iodide (BChI) were used as the substrate in the reaction, 5'dithiobis-2 -nitrobenzoic acid (DTNB) was used as Ellman's reagent were purchased from Sigma - Aldrich. The absorbance of the reaction mixture was measured at 412 nm three times within 5 min of the start of the reaction on a Thermo Fisher Scientific Multiskan Go Finland, and the results are reported as mean  $\pm$  standard deviation. Activity (%) was plotted to determine the inhibitory effects of *V. insulare* extracts on AChE and BChE. IC<sub>50</sub> values were obtained by activity (%) versus compound plots.

### 2.3. The Application of Extracts to Cell Lines

Human colon cancer cell line (HT-29), human breast cancer cell line (MCF-7), and healthy mouse fibroblast cell line (L-929) were used. Dulbecco's Modified Eagle Medium (DMEM) was used as the medium for these cell lines. *V. insulare* extracts were prepared in the DMEM at concentrations of 0.1, 0.2 and 0.5 mg/mL, and the 3-(4,5-dimethyl-thiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) test was applied to determine their cytotoxicity [13]. For the

MTT test,  $5 \times 10^3$  cells were seeded in 100  $\mu\text{L}$  of the medium in each well of 96 well plates. The inoculated cells were incubated in a 5%  $\text{CO}_2$  incubator for 24-48 hours. 100  $\mu\text{l}$  of the diluted extracts in the medium was added to the cell lines. Only 100  $\mu\text{l}$  of the medium was added to the cells in the control wells. Samples were incubated for 24 hours. After incubation, the medium in the wells was removed with the help of a vacuum pump. 10  $\mu\text{l}$  of MTT solution and 90  $\mu\text{l}$  of medium were added to each well and placed in a  $\text{CO}_2$  incubator at  $37^\circ\text{C}$  for 4 hours. After 4 hours, the medium containing MTT was removed from the 96 well plate. 100  $\mu\text{l}$  of Dimethylsulfoxide (DMSO) was added to each well, and their optical density (OD) was measured with a microplate reader (Thermo Fisher Scientific Multiskan Go, Finland) at 540 nm wavelength. A cell line medium without a sample was used as the control group. The average absorbance values obtained by reading the control wells was taken, and this value was accepted as 100% viable cells. The % inhibition rates of the cells were calculated with the help of the following formula.

$$\% \text{ inhibition} = 1 - \left( \frac{\text{OD}_{\text{Sample}}}{\text{OD}_{\text{Control}}} \right) \times 100$$

## 2.4. Statistical Analysis

The % inhibition results of the cell culture and the cholinesterase inhibitory activity were compared among themselves using the t-test following One Way ANOVA. Those with  $p < 0.05$  were statistically significant, and the statistical significance level was indicated with the symbol “\*”. In this respect,  $P < 0.05$  (significant); \*,  $P < 0.01$  (very significant); \*\*,  $P < 0.001$  and  $P < 0.0001$  (extremely significant); \*\*\*-\*\*\*\* and  $P > 0.05$  (not significant), ns.

## 3. Results and Discussion

### 3.1. Anticholinergic Activity

Today, the inhibition of cholinesterases has emerged as an encouraging approach to alleviate the symptoms of Alzheimer's disease. Therefore, inhibition of AChE and BChE enzymes is one of the main goals of researchers. Various cholinesterase inhibitors such as galantamine, tacrine, and rivastigmine are produced synthetically. However, these drugs have side effects such as nausea, diarrhea, dizziness, and vomiting [14]. For these reasons, the discovery of natural and safe enzyme inhibitors has become inevitable. Therefore, the enzyme inhibition properties of *V. insulare* extracts tested against AChE and BChE

enzymes were examined by comparing them with galantamine (Table 1 and Figure 1).

**Table 1.** AChE/BChE inhibition assays for LEtOH (ethanolic leaf), LPW (aqueous leaf pure), REtOH (ethanolic root) and RPW (aqueous root pure) extracts of *V. insulare*

Extracts	AChE inhibition IC <sub>50</sub>	BChE inhibition IC <sub>50</sub>
LEtOH	0.013±0.001	0.042±0.002
LPW	0.014±0.002	0.049±0.003
REtOH	0.038±0.002	0.046±0.002
RPW	0.044±0.005	0.052±0.003
*Galantamine	0.078±0.006	0.117±0.010

\*Galantamine was used as a positive control for Cholinesterase enzymes and determined as  $\mu\text{M}$  levels. Cholinesterase inhibitory activity of the extracts was tested against AChE and BChE at 500  $\mu\text{g}/\text{mL}$  concentration.

All extracts of *V. insulare* exhibited potent AChE and BChE inhibitory activity at 500  $\mu\text{g}/\text{mL}$ . In addition, these extracts showed a better AChE and BChE inhibitory effect than galantamine, a cholinesterase inhibitor used as a therapeutic agent in the treatment of Alzheimer Demands. Galantamine which is used as standard cholinesterase inhibitor was found to have IC<sub>50</sub> values of  $0.078 \pm 0.006 \mu\text{mol}/\text{L}$  and  $0.117 \pm 0.010 \mu\text{mol}/\text{L}$ , respectively, against AChE and BChE enzymes. LEtOH extract of *V. insulare* showed a stronger inhibitory effect for both enzymes than galantamine and other extracts. When the extracts and galantamine activities were compared, it was determined that the extracts showed a highly significant difference from galantamine in general ( $P < 0.0001$ ). It has also been reported that many *Verbascum* species inhibit cholinesterase enzymes. For example, in the study of Angeloni et al., aqueous and methanolic extracts of *Verbascum bombyciferum* were obtained using different techniques such as HAE (homogenizer assisted), MAC (maceration), and infusion was found to exhibit anticholinesterase effects. They found that especially water extracts (maceration:  $0.30 \pm 0.08 \text{ mg GALAE}/\text{g}$  and homogenizer assisted extraction:  $0.28 \pm 0.02 \text{ mg GALAE}/\text{g}$ ) were more weaker inhibitors of AChE when compared to methanolic extracts (homogenizer assisted extraction:  $1.89 \pm 0.15 \text{ mg GALAE}/\text{g}$  and maceration:  $2.02 \pm 0.20 \text{ mg GALAE}/\text{g}$ ) [5]. It was found that aqueous and methanolic extracts of *V. oocarpum* showed an anticholinesterase effect, and the inhibition range varied between  $66.7 \pm 1.2$  and  $28.8 \pm 0.5$  at 200  $\mu\text{g}/\text{mL}$ . As a result, it was determined



acetate extract showed a stronger cytotoxic effect against cancerous cell lines [20]. In a different study, the cytotoxicity of *V. sinaiticum* Benth flower extracts prepared with different solvents against mammary adenocarcinoma MCF-7 was examined, and it was seen that ethanol extract showed the best cytotoxicity [6]. It was also reported that the hydroethanolic leaf extract of the same plant showed the best cytotoxic effect against HepG2 and MRC-5 cell lines [22]. The cytotoxic effect of methanol and hydromethanol extracts obtained from *V. calvum* were compared, and it was reported that the methanol extract displayed important antiproliferative effects against A-549 lung cancer cell line [23]. The plant extracts obtained using different solvents consist of different compounds.

For this reason, it has been determined that the extracts exhibit different anticancer activities in studies. According to the results of this study, it was determined that ethanol extract showed a better antiproliferative effect than pure water extract. This study supports the view that the cytotoxic effect may be due to the compounds contained in the extract, as in previous studies.

#### 4. Conclusion and Suggestions

It is known that the damage caused by oxidative stress in cells and tissues may play a role in the development of many diseases such as Alzheimer's and cancer in the organism. In the key enzyme inhibition theory, which is effective for the management of neurological diseases such as Alzheimer's, inhibition of these enzymes can alleviate existing symptoms. For this purpose, the presence of natural inhibitors for cholinesterase inhibition is very important. In this study, anticancer activity of *V.insulare*, a species belonging to the genus *Verbascum*, against human colon cancer cell line (HT-29), human breast cancer cell line (MCF-7), and healthy mouse fibroblast cell line (L-929), and enzyme (AChE, BChE) inhibitory effects were investigated for the first time. It was determined that the ethanol extract from the extracts obtained by using different plant parts and solvents showed better anticholinesterase activity and antiproliferative properties than the pure water extract. In addition, leaf ethanol extract showed the best activity for both biological activity tests. This is thought to be due to the fact that the extracts prepared in different parts of the plant and in different solvents contain different compounds. It is anticipated that future studies will isolate and purify the active compounds and investigate their effects on a wider variety of cancer cells and enzymes. However, *in vivo* tests, namely more studies, are needed to explore the

full mode of action of the active compounds in herbal medicine development and to test the safety of the extracts used here.

#### Contributions of the authors

Corresponding author: Data curation, writing (original draft, review & editing), investigation, visualization methodology, *in vitro* studies and interpretation of experimental results.

Coauthor: Methodology, investigation, visualization, literature review, *in vitro* studies, and interpretation of experimental results.

#### Conflict of Interest Statement

There is no conflict of interest between the authors.

#### Statement of Research and Publication Ethics

Research and publication ethics compiled within the study.

## References

- [1] S.-Y. Pan *et al.*, "New perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics," *Evidence-Based Complementary and Alternative Medicine*, vol. 2013, 2013.
- [2] Y. Alan and N. Yilmaz, "Phenolic substance contents and biological activities of verbascum insulare boiss. and heldr. Extracts," *J steroids*, vol. 33, p. 35, 2019.
- [3] I. Süntar, I. I. Tatlı, E. K. Akkol, H. Keleş, Ç. Kahraman, and Z. Akdemir, "An ethnopharmacological study on Verbascum species: from conventional wound healing use to scientific verification," *Journal of ethnopharmacology*, vol. 132, no. 2, pp. 408-413, 2010.
- [4] H. A. E. H. El Gizawy, M. A. Hussein, and E. Abdel-Sattar, "Biological activities, isolated compounds and HPLC profile of Verbascum nubicum," *Pharmaceutical biology*, vol. 57, no. 1, pp. 485-497, 2019.
- [5] S. Angeloni *et al.*, "An insight into Verbascum bombyciferum extracts: Different extraction methodologies, biological abilities and chemical profiles," *Industrial Crops and Products*, vol. 161, p. 113201, 2021.
- [6] W. H. Talib and A. M. Mahasneh, "Antiproliferative activity of plant extracts used against cancer in traditional medicine," *Scientia pharmaceutica*, vol. 78, no. 1, pp. 33-46, 2010.
- [7] Y.-L. Zhao *et al.*, "Isolation of chemical constituents from the aerial parts of Verbascum thapsus and their antiangiogenic and antiproliferative activities," *Archives of pharmacal research*, vol. 34, no. 5, pp. 703-707, 2011.
- [8] J. A. Dumas and P. A. Newhouse, "The cholinergic hypothesis of cognitive aging revisited again: cholinergic functional compensation," *Pharmacology Biochemistry and Behavior*, vol. 99, no. 2, pp. 254-261, 2011.
- [9] J. E. Lee, D. Kim, and J. H. Lee, "Association between alzheimer's disease and cancer risk in South Korea: an 11-year nationwide population-based study," *Dementia and neurocognitive disorders*, vol. 17, no. 4, pp. 137-147, 2018.
- [10] A. F. Yücel, Ö. Kemik, A. S. Kemik, S. Purisa, and İ. S. Tüzün, "Relationship between the levels of oxidative stress in mesenteric and peripheral serum and clinicopathological variables in colorectal cancer," *Balkan Medical Journal*, vol. 2012, no. 2, pp. 144-147, 2012.
- [11] G. L. Ellman, K. D. Courtney, V. Andres Jr, and R. M. Featherstone, "A new and rapid colorimetric determination of acetylcholinesterase activity," *Biochemical pharmacology*, vol. 7, no. 2, pp. 88-95, 1961.
- [12] Z. Almaz, A. Oztekin, A. Tan, and H. Ozdemir, "Biological evaluation and molecular docking studies of 4-aminobenzohydrazide derivatives as cholinesterase inhibitors," *Journal of Molecular Structure*, p. 130918, 2021.
- [13] M. E. Alkış, Ü. Keleştemür, Y. Alan, N. Turan, and K. Buldurun, "Cobalt and ruthenium complexes with pyrimidine based schiff base: Synthesis, characterization, anticancer activities and electrochemotherapy efficiency," *Journal of Molecular Structure*, vol. 1226, p. 129402, 2021.
- [14] N. Martins and I. C. Ferreira, "Neurocognitive improvement through plant food bioactives: A particular approach to Alzheimer's disease," in *Food Bioactives*: Springer, 2017, pp. 267-298.
- [15] S. Kanbolat *et al.*, "Antioxidant, antimicrobial, cytotoxic, anticholinesterase, antityrosinase activities and characterisation of volatile compounds of Verbascum oocarpum by SPME and GC-FID/MS," *Journal of Pharmaceutical Research International*, pp. 1-12, 2018.
- [16] C. Kahraman, I. I. Tatlı, I. E. Orhan, and Z. S. Akdemir, "Cholinesterase inhibitory and antioxidant properties of Verbascum mucronatum Lam. and its secondary metabolites," *Zeitschrift für Naturforschung C*, vol. 65, no. 11-12, pp. 667-674, 2010.
- [17] G. Zengin *et al.*, "Evaluation of chemical constituents and biological properties of two endemic Verbascum species," *Process Biochemistry*, vol. 108, pp. 110-120, 2021.
- [18] J. Iqbal *et al.*, "Plant-derived anticancer agents: A green anticancer approach," *Asian Pacific Journal of Tropical Biomedicine*, vol. 7, no. 12, pp. 1129-1150, 2017.
- [19] G. M. Cragg and D. J. Newman, "Plants as a source of anti-cancer agents," *Journal of ethnopharmacology*, vol. 100, no. 1-2, pp. 72-79, 2005.
- [20] İ. İ. TATLI and Z. S. Akdemir, "Cytotoxic activity on some Verbascum species growing in Turkey," *Hacettepe University Journal of the Faculty of Pharmacy*, no. 2, pp. 77-86, 2006.

- [21] I. A. Iliescu *et al.*, "Verbascum nigrum: cytotoxicity evaluation in A431 epidermoid carcinoma cells and untargeted LC-HR-MS/MS metabolite profiling," *Chemistry & Biodiversity*, vol. 17, no. 12, p. e2000644, 2020.
- [22] J. Tauchen *et al.*, "In vitro antioxidant and anti-proliferative activity of Ethiopian medicinal plant extracts," *Industrial Crops and Products*, vol. 74, pp. 671-679, 2015.
- [23] H. I. M Amin, F. H. Hussain, G. Gilardoni, Z. M. Thu, M. Clericuzio, and G. Vidari, "Phytochemistry of Verbascum Species Growing in Iraqi Kurdistan and Bioactive Iridoids from the Flowers of Verbascum calvum," *Plants*, vol. 9, no. 9, p. 1066, 2020.